

A wonderful Bio-control agent of 21st Century

Bio-Abhijat Liq.

Pseudomonas fluorescens

(Spore Count (CFU) — 2×10^{10} /gm)

Pseudomonas fluorescens controls root & stem rots caused by **Schlerotina** and **Rhizoctonia**, wilting caused by **Fusarium**, **Erwinia**, damping off caused by **Pythium**, blights / leaf spots caused by **Alternaria**, **Ascochyta**, **Cercospora**, **Macrophomina**, **Myrothecium** & **Ramularia**, downy mildew, powdery mildew and Fungal diseases. When applied in the soil, the bacteria move towards the plant roots and grow profusely over the root surface. It then enters the vascular streams,

reach the leaves and act as a systemic biocontrol agent against diseases. The bacteria produces antibiotics. Once it grows over the root surface, it acts as a shield against the invading plant pathogens. Further, it secretes several plant growth substances, and these gibberelin - like compounds contribute to vigorous growth of the plant. *Pseudomonas* also helps in remediation of soil from harmful effects of heavy metals & herbicides.

Advantages :

- Protects crops from soil borne diseases.
- Protects crops from diseases throughout the crop period.
- Do not leave harmful residues.
- It is target specific and do not destroy beneficial organisms.
- Environmental friendly & have no adverse effect on soil or plants or humans or animals.

Dosage & Direction :

1. **Seed or Seedling root treatment** : Mix Bio-Abhijat Liq. at the rate 2 ml in 1 Ltr. of water. Soak the seeds or root of seedlings for 30 minutes, before sowing or planting.
2. **Soil Application** : Mix 200 ml of Bio-Abhijat Liq. with required quantity (300 Kg/Acre) of organic manure or Farm Yard Manure (FYM) or vermicompost for covering one Acre area. After mixing leave the mix in a heap at a covered, dry and cool place for 24 hours and then apply in the soil. This soil application can be done 3 days before or within 10 to 15 days after sowing or planting.
3. **Foliar application** : Mix Bio-Abhijat Liq. at the rate 1 ml in 1 Ltr. of water, with the required quantity of water for 1 Acre or mix 200 ml in 200 Ltr. of water and spray evenly on the foliage. Also drench the soil near the root zone of the plant. The first spray can be done between 30 to 45 days after sowing or planting. The spray can be repeated once 20 to 30 days depending upon the persistence of the disease.

Dosage :

Seed application : 1 ml / kg seed.	Seedling application : 2 ml / l water	Soil application : 200 ml / Acre with 300 kg FYM or Compost	Foliar application : 1 ml / l water
----------------------------------------------	-------------------------------------------------	--------------------------------------------------------------------------	-----------------------------------------------

Certified by :
biocertTM
International Pvt. Ltd.

NIVSHAKTI
BIDENERGY PVT LTD
In Harmony with Nature
An ISO 9001:2008 Company
Formerly known as
SHAKTI BIOTECH

Junaid Manzil, 2nd Floor,
7B, Ahiripukur Road, Kolkata - 700 019
Ph : +91 33 2287 2244 / 2400
E : support@shaktibiotech.in
W : www.shaktibiotech.in

Be OrganicTM

MEMBER



CSR PARTNER



Warranty : The information supplied in this leaflet is true to our knowledge and based on our experience. However, no liability is accepted for any insufficiency or omission neither any responsibility is assumed for any procedural deviation in the actual working place.